

MICROCOPY RESOLUTION TEST CHART

2

NPS-63-84-006

NAVAL POSTGRADUATE SCHOOL Monterey, California

AD-A146 521



Monthly and Seasonal Climatology over the
Global Tropics and Subtropics
for the Decade 1973 to 1983

Volume I. 200 mb winds

by

James S. Boyle and C.-P. Chang

August 1984

Technical Report August 1983 - August 1984

Approved for public release; distribution unlimited.

Prepared for: National Oceanic and Atmospheric Administration
Washington, D.C. 20233

84 10 01 106

DTIC ELECTED
OCT 10 1984
S E

DTIC FILE COPY

NAVAL POSTGRADUATE SCHOOL
Monterey, California

Commodore R. H. Shumaker
Superintendent

D. A. Schradly
Provost

The work reported herein was supported by the National Oceanic and Atmospheric Administration, under Research Contract NA83MAG03828.

This report was prepared by:

James S. Boyle
JAMES S. BOYLE
Adjunct Professor of Meteorology

C P Chang
C.-P. CHANG
Professor of Meteorology

Publication of the report does not constitute approval of the sponsor for the findings or conclusions. It is published for information and for the exchange and stimulation of ideas.

Reviewed by:

Robert J. Renard
ROBERT J. RENARD
Chairman
Department of Meteorology

C P Chang
C.-P. CHANG
Professor of Meteorology
Principal Investigator

Released by:

John N. Dyer
JOHN N. DYER
Dean of Science and Engineering

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS-63-84-006	2. GOVT ACCESSION NO. AD-A146521	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Monthly and Seasonal Climatology over the Global Tropics and Subtropics for the Decade 1973 to 1983. Volume I. 200 mb winds.		5. TYPE OF REPORT & PERIOD COVERED Technical Report for August 1983 - August 1984
7. AUTHOR(s) J. S. Boyle and C.-P. Chang		6. PERFORMING ORG. REPORT NUMBER NOAA Research Contract NA63AAGC03828
8. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93943		9. CONTRACT OR GRANT NUMBER(s) NOAA Research Contract NA63AAGC03828
10. CONTROLLING OFFICE NAME AND ADDRESS Climate Analysis Center/NMC National Oceanic and Atmospheric Administration Washington, D.C. 20233		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 1984
		13. NUMBER OF PAGES 172
		14. SECURITY CLASS. (of this report) UNCLASSIFIED
15. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		16. SECURITY CLASSIFICATION/DOWNGRADING SCHEDULE

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Interannual Variations
Tropical Climatology
Winter Circulations

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

This atlas of the 200 mb circulation field contains northern winter monthly and seasonal mean wind analyses, velocity potential and streamfunction from 40S to 60N over a global belt for the decade 1973 through 1983. In addition, the deviations of the individual annual seasonal and monthly means from their respective ten year means are presented for the same variables. The basic wind data used are the operational Global Band Analyses (GBA) of the United States Navy's Fleet Numerical Oceanography Center (FNOC). The analyses exhibit many interesting features, interannual variations and may shed some light on the understanding of the El Niño/Southern Oscillation phenomena.

DD FORM 1 JAN 78 1473 EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-010-0001

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Contents

Abstract _____ 1

1. Introduction _____ 2

2. Data Sources, Analyses and Computations _____ 3

 2.1 Global Band Analysis _____ 3

 2.2 Computation of Streamfunction and Velocity Potential _____ 4

3. Discussion _____ 5

 3.1 Introduction _____ 5

 3.2 El Nino/Southern Oscillation Events _____ 5

 3.3 Decade Overview _____ 6

References _____ 7

Figures _____ 8

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



ABSTRACT

This atlas of the 200 mb circulation field contains northern winter monthly and seasonal mean wind analyses, velocity potential and streamfunction from 40S to 60N over a global belt for the decade 1973 through 1983. In addition, the deviations of the individual annual seasonal and monthly means from their respective ten year means are presented for the same variables. The basic wind data used are the operational Global Band Analyses (GBA) of the United States Navy's Fleet Numerical Oceanography Center (FNOOC). The analyses exhibit many interesting features, interannual variations and may shed some light on the understanding of the El Nino/Southern Oscillation phenomena.

1. INTRODUCTION

This atlas depicts the wintertime (December, January, February) seasonal and monthly mean atmospheric motion field at 200 mb for the decade 1973/1974 to 1982/1983. The charts display wind vectors, isotachs, velocity potential and streamfunction from 60N to 40S at 200 mb. In addition to the ten-year seasonal and monthly means and the individual annual seasonal and monthly averages, the deviations of the individual seasonal and monthly means from their ten-year averages are also presented. The seasonal calculations are based on the months of December, January, and February. To aid in depicting the onset of the winter monsoon circulations the monthly charts include the November as well as December, January, and February. The data used to derive these motion fields are the Operational Global Band Analysis (GBA) of the United States Navy's Fleet Numerical Oceanography Center (FNOC). The procedures used in producing these analyses are described in section 2.

For reference seasonal and monthly means and deviations therefrom of the outgoing longwave radiation (OLR) as observed from the NOAA polar orbiting satellites for the period from 1974/1975 to 1982/1983 are available in volume II of this report.

The motion fields for the decade of winters from 1973/74 to 1982/83 are of interest since the data cover a period not hitherto examined in other collections of data. Other works such as Oort (1983) and Krishnamurti et al. (1983) have presented detailed analyses of the decade prior to 1973. The 1973-1983 decade contains two El Nino/Southern Oscillation (ENSO) events, one occurring in 1976/77 and the other in 1982/83, the latter event is the most intense ENSO events yet observed. Also, the analyses presented here allow the FGGE winter to be placed in a longer term perspective since the FGGE/MMONEX experiment took place in the midst of the decade.

2. DATA SOURCES, ANALYSES AND COMPUTATIONS

2.1 GLOBAL BAND ANALYSIS

The wind data set used in this work are the operational analyses of the Global Band Analyses (GBA) of FNOC. These data are produced four times daily by objective procedures on a mercator grid which extends from 60N to 40S. The use of the mercator secant projection results in a change in the actual distance between grid points from 140 km at 60N to a maximum value of 280 km at the equator. The objective scheme is designed to take advantage of all the reports in the operational data base, surface synoptic, aircraft, pilot balloons, rawinsonde and satellite data.

The analysis is performed every six hours for the surface, 700, 400, 250 and 200 mb levels. The first guess field used as input for the objective analysis is the six-hour persistence field. The operational approach is to first interpolate the irregularly spaced data to grid points using a successive corrections technique based on Cressman's (1959) method. The successive corrections method takes several scans through the data reducing the scan radius on each successive scan. Analyses are performed of both wind and temperature by this method. These fields are then adjusted by a set of numerical variational analysis (NVA) equations which have incorporated the dynamical constraints of the momentum equations with friction included in the surface layer (Lewis and Grayson, 1972). Temperature and wind fields are adjusted subject to mutual constraints on the fields. However, the 200 mb wind data used in this atlas serve only as a boundary condition for NVA and are not subject to the adjustment.

2.2 COMPUTATION OF STREAMFUNCTION AND VELOCITY POTENTIAL

The streamfunction (ψ) and velocity potential (χ) were computed from the following equations:

$$\nabla^2 \psi = \zeta \quad (1)$$

$$\nabla^2 \chi = \delta \quad (2)$$

where:

$$\zeta \text{ is the relative vorticity} = \partial v / \partial y - \partial u / \partial x$$

and

$$\delta \text{ is the divergence} = \partial u / \partial x + \partial v / \partial y.$$

Both ζ and δ were computed using centered differences on the GBA mercator grid.

Equation 2 was solved using the boundary condition that $\chi = 0$ at the north and south boundaries which are 60N and 40S respectively. The method used to compute ψ was essentially method II of Shukla and Seha (1974). This technique uses the previously computed values of χ to formulate boundary conditions for ψ . The values of ψ above 50N are not displayed because of obvious difficulties that the solution for ψ encountered on this boundary. It was felt that displaying these results detracted from the overall presentation. Although the boundary condition described here produced the best results of the many different methods tried, the location of the boundary in the meteorologically active region at 60N proved to be troublesome. The depiction of the divergence field appears reasonable away from the boundary. Comparison with the global fields produced by the National Meteorological Center (NMC) for the years since the NMC global product has become available indicate that the effect of the boundary condition on χ is not significant between 40N and 30S. Thus the ψ and χ fields in the equatorial regions are sufficiently removed from the boundaries that we can assume that these values are not unduly affected by the choice of boundary conditions.

3. DISCUSSION

3.1 INTRODUCTION

This discussion is not intended to be a comprehensive review of the tremendous amount of data contained in the accompanying figures. Rather the intent is to describe in general terms some of the more obvious general features of interest.

3.2 EL NINO/SOUTHERN OSCILLATION EVENTS

A perusal of the charts of the seasonal means of 200 mb χ clearly show that the dominant center of interannual variability is located in the central Pacific. These fluctuations can be associated with changes driven by the Southern Oscillation.

There are a great many studies in the literature about the global scale aspects of the ENSO events. The particular collection of charts presented here depicts the wintertime 200 mb motion fields for two such events occurring in 1976/77 and 1982/83. Liebmann and Hartmann (1982) describe the anomalous (based on a three year average) patterns of OLR for the 1976/77 event. They show the increase in convection activity over Indonesia shifts eastward after the sea-surface temperature (SST) in the east Pacific becomes anomalously warm. The South Pacific convergence zone (SPCZ) intensifies and moves eastward but does not reach its maximum intensity until more than a year after the SST increases occur. Lau et al. (1983) consider the anomalous OLR, ψ and χ patterns for both the 76/77 and 82/83 ENSO events using the same data as presented here (OLR data is presented in volume II). They concluded that the circulation changes associated with each individual ENSO event can be quite different, but that the anomalous OLR patterns and circulation

features within each event were consistent. They showed that prior to the ENSO of 1976/77 the equatorial central Pacific convection was strongly suppressed for at least a two year period prior to December 1975. The seasonal data of 1976/77 and 1977/78 evince a dramatic reversal in the X pattern in the eastern Pacific.

3.3 DECADE OVERVIEW

If one examines the seasonal means of X and OLR it becomes apparent that there is a shift in the equatorial Pacific about the middle of the decade. Figure I is a time series of the deviations from the ten-year mean of the winter season X field averaged over the region from 175E to 140W and 10N to 10S. This region, located in the mid-Pacific, is one of large inter-seasonal variability. The figure indicates two regimes, one going from 1973/75 to 1976/77, the other from 1977/78 to 1982/83. The former has relatively low values of X, the latter relatively high values. This can be interpreted as an increase in the upper-level divergence in this region in the latter part of the decade. Figure I also contains a time series of the deviations from the nine-year mean of the winter season OLR (winter 1973/74 not available) averaged over the same region as the X values. Negative values indicate relatively decreased OLR which can be construed to mean decreased convection in the tropics. Figure I demonstrates a strong negative correlation, which implies that the independently derived data sets are consistent on the seasonal time scale. Note that the OLR values are plotted on an inverse scale from the X graph. This also indicates that the FGCE winter (1978/79) is more characteristic of the regime in the latter half of the decade and care must be taken extrapolating inferences drawn from the FGCE data to other years.

REFERENCES

- Cressman, G. P., 1959: An operational objective analysis system. Mon. Wea. Rev., 87, 367-374.
- Gruber, A. and J. S. Winston, 1978: Earth-atmosphere radiative heating based on NOAA scanning radiometer measurements. Bull. Amer. Meteor. Soc., 59, 1570-1573.
- Krishnamurti, T. N., H.-L. Pan, R. Pasch and D. Subrahmanyam, 1983: Interannual variability of the tropical motion field. FSU Report No. 83-3, Department of Meteorology, Florida State University, Tallahassee, Florida 32306.
- Lau, K.-M., J. S. Boyle and C.-P. Chang, 1984: Anomalous diabatic heating and large scale circulation in the tropics during the 1976-77 and 1982-83 ENSO's. Tropical Ocean - Atmosphere Newsletter, 24, 12-13.
- Lau, K.-M. and P. H. Chan, 1983: Short-term climate variability and atmospheric teleconnections from satellite observed outgoing longwave radiation. Part I: Simultaneous relationships. J. Atmos. Sci., 40, 2735-2750.
- Lewis, J. M. and T. H. Grayson, 1972: The adjustment of surface wind and pressure by Sasaki's variational matching technique. J. Appl. Meteor., 11, 586-597.
- Liebmann, B. and D. L. Hartmann, 1982: Interannual variation of outgoing IR associated with tropical circulation changes during 1974-1978. J. Atmos. Sci., 39, 1153-1162.
- Oort, A. H., 1983: Global atmospheric circulation statistics, 1958-1973. NOAA Professional Paper 14. U.S. Department of Commerce (available from the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
- Shukla, J. and K. R. Saha, 1974: Computation of non-divergent streamfunction and irrotational velocity potential from the observed winds. Mon. Wea. Rev., 102, 419-425.

FIGURES

Figure I

Time series of the deviations of the seasonal means of χ and OLR from their respective ten- and nine- year seasonal means. The values are areal averages over the region bounded by 10N to 10S and 175E to 140W. The χ values are represented by the solid line, the OLR values by the dashed line. Note that the OLR ordinate scale (on the left hand side of the graph) is reversed from that of the χ ordinate. The ten-year mean value of χ for the region is $4.6 \times 10^{-6} \text{ m}^{-1} \text{ s}^{-2}$. The nine-year mean of OLR for the region is 234 Wm^2 .

Figures A1 - A3

Ten winter season (1973/74 - 1982/83) mean circulation fields at 200 mb. Variables displayed are streamfunction, velocity potential, wind vectors and isotachs. Contour interval for the streamfunction is $1.0 \times 10^7 \text{ m}^2 \text{ s}^{-1}$, for velocity potential it is $1.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, and for isotachs it is 10 ms^{-1} . The vector scale is given in the upper right portion of the wind vector isotach plots. The grid intervals of the FNOC Global Band Analysis mercator grid are shown on the left hand side and bottom of each figure. The longitude grid is marked every 30° from the Greenwich meridian on the extreme left and the latitude is marked every 10° from 40S at the bottom of the figure.

Figures B1 - B30

Individual winter season mean and deviation circulation fields at 200 mb, for the winters from 1973/74 to 1982/83. The deviations are differences from the ten year seasonal mean (Figs. A1 and A3). The figures are marked with the year corresponding to the year of the January and February of the winter. Variables displayed are the streamfunction, velocity potential, wind vectors and isotachs. For the mean fields the contour interval for the streamfunction is $1.0 \times 10^7 \text{ m}^2 \text{ s}^{-1}$, for velocity potential it is $1.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, and for isotachs it is 20 ms^{-1} . For the deviation fields the contour interval for the streamfunction is $2.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, for velocity potential it is $1.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, and for isotachs it is 10 ms^{-1} . On the deviation plots contours with negative values are dashed, those with positive values and the zero line are solid. The vector scale is given in the upper right portion of the wind vector isotach plots.

Figure C1 - C12

Ten-year (1973 - 1983) monthly mean circulation fields at 200 mb. The months displayed are November, December, January, and February. Variables displayed are streamfunction, velocity potential, wind vectors and isotachs. Contour interval for the streamfunction is $1.0 \times 10^7 \text{ m}^2 \text{ s}^{-1}$, for velocity potential it is $1.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, and for isotachs it is 20 ms^{-1} . The vector scale is given in the upper right portion of the wind vector isotach plots.

Figures D1 - D120

Individual monthly mean and deviation circulation fields at 200 mb, for the months from November 1973 to February 1983. The deviations are differences from the ten-year monthly means (Figs. C1 and C12). The months displayed are November, December, January, and February. Variables displayed are the streamfunction, velocity potential, wind vectors and isotachs. For the mean fields the contour interval for the streamfunction is $1.0 \times 10^7 \text{ m}^2 \text{ s}^{-1}$, for velocity potential it is $1.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, and for isotachs it is 20 ms^{-1} . For the deviation fields the contour interval for the streamfunction is $1.0 \times 10^7 \text{ m}^2 \text{ s}^{-1}$, for velocity potential it is $1.0 \times 10^6 \text{ m}^2 \text{ s}^{-1}$, and for isotachs it is 20 ms^{-1} . On the deviation plots contours with negative values are dashed, those with positive values and the zero line are solid. The vector scale is given in the upper right portion of the wind vector isotach plots.





Figure 1

STREAM FUNCTION 200MB WIN AVG


A1



OBSERVED WIND 200MB WIN AVG


A2



SCALE = 20




VELOCITY POTENTIAL 200MB WIN AVG

A3







STREAM FUNCTION 200MB WIN 1974




STREAM FUNCTION 200MB WIN 1974



VELOCITY POTENTIAL 200MB WIN 1974



VELOCITY POTENTIAL 200MB WIN 1974




OBSERVED WIND 200MB WIN 1975

SCALE = 20




OBSERVED WIND 200MB WIN 1975


SCALE = 10




STREAM FUNCTION 200MB WIN 1975




STREAM FUNCTION 200MB WIN 1975





VELOCITY POTENTIAL 200MB WIN 1976




VELOCITY POTENTIAL 200MB WIN 1976







STREAM FUNCTION 200MB WIN 1976




STREAM FUNCTION 200MB WIN 1976



VELOCITY POTENTIAL 200MB WIN 1976




VELOCITY POTENTIAL 200MB WIN 1976




OBSERVED WIND 200MB WIN 1977

SCALE = 20




OBSERVED WIND 200MB WIN 1977


SCALE = 10




STREAM FUNCTION 200MB WIN 1977




STREAM FUNCTION 200MB WIN 1977



VELOCITY POTENTIAL 200MB WIN 1977




VELOCITY POTENTIAL 200MB WIN 1977




OBSERVED WIND 200MB WIN 1978

SCALE = 20




OBSERVED WIND 200MB WIN 1978


SCALE = 10




STREAM FUNCTION 200MB WIN 1978




STREAM FUNCTION 200MB WIN 1978



VELOCITY POTENTIAL 200MB WIN 1978



VELOCITY POTENTIAL 200MB WIN 1978




OBSERVED WIND 200MB WIN 1979

SCALE = 20




OBSERVED WIND 200MB WIN 1979


SCALE = 10




STREAM FUNCTION 200MB WIN 1979




STREAM FUNCTION 200MB WIN 1979



VELOCITY POTENTIAL 200MB WIN 1979




VELOCITY POTENTIAL 200MB WIN 1979




OBSERVED WIND 200MB WIN 1980

SCALE = 20




OBSERVED WIND 200MB WIN 1980


SCALE = 10




STREAM FUNCTION 200MB WIN 1980




STREAM FUNCTION 200MB WIN 1980





VELOCITY POTENTIAL 200MB WIN 1980




VELOCITY POTENTIAL 200MB WIN 1980







STREAM FUNCTION 200MB WIN 1981




STREAM FUNCTION 200MB WIN 1981



VELOCITY POTENTIAL 200MB WIN 1981




VELOCITY POTENTIAL 200MB WIN 1981




OBSERVED WIND 200MB WIN 1982

SCALE = 20




OBSERVED WIND 200MB WIN 1982


SCALE = 10




STREAM FUNCTION 200MB WIN 1982




STREAM FUNCTION 200MB WIN 1982



VELOCITY POTENTIAL 200MB WIN 1982




VELOCITY POTENTIAL 200MB WIN 1982




OBSERVED WIND 200MB WIN 1983

SCALE = 20




OBSERVED WIND 200MB WIN 1983


SCALE = 10




STREAM FUNCTION 200MB WIN 1983




STREAM FUNCTION 200MB WIN 1983



VELOCITY POTENTIAL 200MB WIN 1983




VELOCITY POTENTIAL 200MB WIN 1983



STREAM FUNCTION 200MB NOV AVG


C1




OBSERVED WIND 200MB NOV AVG


C2

SCALE = 20






STREAM FUNCTION 200MB DEC AVG




OBSERVED WIND 200MB DEC AVG



SCALE = 20


VELOCITY POTENTIAL 200MB DEC AVG



C6

STREAM FUNCTION 200MB JAN AVG


C7



OBSERVED WIND 200MB JAN AVG


C8

SCALE = 20




VELOCITY POTENTIAL 200MB JAN AVG

C9



C10


STREAM FUNCTION 200MB FEB AVG



C11


SCALE = 20

OBSERVED WIND 200MB FEB AVG




VELOCITY POTENTIAL 200MB FEB AVG

C12




SCALE = 20

OBSERVED WIND 200MB NOV 1973




SCALE = 20


OBSERVED WIND 200MB NOV 1973




STREAM FUNCTION 200MB NOV 1973




STREAM FUNCTION 200MB NOV 1973



VELOCITY POTENTIAL 200MB NOV 1973




VELOCITY POTENTIAL 200MB NOV 1973




OBSERVED WIND 200MB DEC 1973

SCALE = 20




OBSERVED WIND 200MB DEC 1973


SCALE = 20




STREAM FUNCTION 200MB DEC 1973




STREAM FUNCTION 200MB DEC 1973



VELOCITY POTENTIAL 200MB DEC 1973




VELOCITY POTENTIAL 200MB DEC 1973




SCALE = 20

OBSERVED WIND 200MB JAN 1974




SCALE = 20


OBSERVED WIND 200MB JAN 1974




STREAM FUNCTION 200MB JAN 1974




STREAM FUNCTION 200MB JAN 1974



VELOCITY POTENTIAL 200MB JAN 1974



VELOCITY POTENTIAL 200MB JAN 1974




OBSERVED WIND 200MB FEB 1974

SCALE = 20




OBSERVED WIND 200MB FEB 1974


SCALE = 20




STREAM FUNCTION 200MB FEB 1974




STREAM FUNCTION 200MB FEB 1974



VELOCITY POTENTIAL 200MB FEB 1974




VELOCITY POTENTIAL 200MB FEB 1974




SCALE = 20

OBSERVED WIND 200MB NOV 1974




SCALE = 20


OBSERVED WIND 200MB NOV 1974




STREAM FUNCTION 200MB NOV 1974




STREAM FUNCTION 200MB NOV 1974



VELOCITY POTENTIAL 200MB NOV 1974




VELOCITY POTENTIAL 200MB NOV 1974




OBSERVED WIND 200MB DEC 1974

SCALE = 20




OBSERVED WIND 200MB DEC 1974


SCALE = 20




STREAM FUNCTION 200MB DEC 1974




STREAM FUNCTION 200MB DEC 1974



VELOCITY POTENTIAL 200MB DEC 1974




VELOCITY POTENTIAL 200MB DEC 1974




OBSERVED WIND 200MB JAN 1976

SCALE = 20




OBSERVED WIND 200MB JAN 1976


SCALE = 20




STREAM FUNCTION 200MB JAN 1976




STREAM FUNCTION 200MB JAN 1976



VELOCITY POTENTIAL 200MB JAN 1975




VELOCITY POTENTIAL 200MB JAN 1975




OBSERVED WIND 200MB FEB 1975

SCALE = 20




OBSERVED WIND 200MB FEB 1975


SCALE = 20







STREAM FUNCTION 200MB FEB 1975




STREAM FUNCTION 200MB FEB 1975









STREAM FUNCTION 200MB NOV 1975




STREAM FUNCTION 200MB NOV 1975



VELOCITY POTENTIAL 200MB NOV 1975




VELOCITY POTENTIAL 200MB NOV 1975




OBSERVED WIND 200MB DEC 1976

SCALE = 20




OBSERVED WIND 200MB DEC 1976


SCALE = 20




STREAM FUNCTION 200MB DEC 1975




STREAM FUNCTION 200MB DEC 1975



VELOCITY POTENTIAL 200MB DEC 1975




VELOCITY POTENTIAL 200MB DEC 1975



OBSERVED WIND 200MB JAN 1976

SCALE = 20




OBSERVED WIND 200MB JAN 1976


SCALE = 20




STREAM FUNCTION 200MB JAN 1976




STREAM FUNCTION 200MB JAN 1976



VELOCITY POTENTIAL 200MB JAN 1976




VELOCITY POTENTIAL 200MB JAN 1976




OBSERVED WIND 200MB FEB 1976

SCALE = 20




OBSERVED WIND 200MB FEB 1976


SCALE = 20




STREAM FUNCTION 200MB FEB 1976




STREAM FUNCTION 200MB FEB 1976



VELOCITY POTENTIAL 200MB FEB 1976




VELOCITY POTENTIAL 200MB FEB 1976




OBSERVED WIND 200MB NOV 1976

SCALE = 20




OBSERVED WIND 200MB NOV 1976


SCALE = 20




STREAM FUNCTION 200MB NOV 1976




STREAM FUNCTION 200MB NOV 1976



VELOCITY POTENTIAL 200MB NOV 1976




VELOCITY POTENTIAL 200MB NOV 1976




OBSERVED WIND 200MB DEC 1976


SCALE = 20



OBSERVED WIND 200MB DEC 1976


SCALE = 20






MICROCOPY RESOLUTION TEST CHART


STREAM FUNCTION 200MB DEC 1976




STREAM FUNCTION 200MB DEC 1976





VELOCITY POTENTIAL 200MB DEC 1976




VELOCITY POTENTIAL 200MB DEC 1976







STREAM FUNCTION 200MB JAN 1977




STREAM FUNCTION 200MB JAN 1977







VELOCITY POTENTIAL 200MB JAN 1977



VELOCITY POTENTIAL 200MB JAN 1977









OBSERVED WIND 200MB NOV 1977

SCALE = 20



OBSERVED WIND 200MB NOV 1977


SCALE = 20




100




STREAM FUNCTION 200MB NOV 1977




STREAM FUNCTION 200MB NOV 1977



VELOCITY POTENTIAL 200MB NOV 1977




VELOCITY POTENTIAL 200MB NOV 1977




OBSERVED WIND 200MB DEC 1977

SCALE = 20




OBSERVED WIND 200MB DEC 1977


SCALE = 20




STREAM FUNCTION 200MB DEC 1977




STREAM FUNCTION 200MB DEC 1977



VELOCITY POTENTIAL 200MB DEC 1977




VELOCITY POTENTIAL 200MB DEC 1977




SCALE = 20

OBSERVED WIND 200MB JAN 1978




SCALE = 20


OBSERVED WIND 200MB JAN 1978




STREAM FUNCTION 200MB JAN 1978




STREAM FUNCTION 200MB JAN 1978



VELOCITY POTENTIAL 200MB JAN 1978




VELOCITY POTENTIAL 200MB JAN 1978




OBSERVED WIND 200MB FEB 1978

SCALE = 20




OBSERVED WIND 200MB FEB 1978


SCALE = 20



VELOCITY POTENTIAL 200MB FEB 1978




VELOCITY POTENTIAL 200MB FEB 1978




OBSERVED WIND 200MB NOV 1978

SCALE = 20




OBSERVED WIND 200MB NOV 1978


SCALE = 20




STREAM FUNCTION 200MB NOV 1978




STREAM FUNCTION 200MB NOV 1978



VELOCITY POTENTIAL 200MB NOV 1978



VELOCITY POTENTIAL 200MB NOV 1978




OBSERVED WIND 200MB DEC 1978

SCALE = 20




OBSERVED WIND 200MB DEC 1978


SCALE = 20




STREAM FUNCTION 200MB DEC 1978




STREAM FUNCTION 200MB DEC 1978



VELOCITY POTENTIAL 200MB DEC 1978




VELOCITY POTENTIAL 200MB DEC 1978




OBSERVED WIND 200MB JAN 1979

SCALE = 20




OBSERVED WIND 200MB JAN 1979


SCALE = 20




STREAM FUNCTION 200MB JAN 1978




STREAM FUNCTION 200MB JAN 1979



VELOCITY POTENTIAL 200MB JAN 1979



VELOCITY POTENTIAL 200MB JAN 1979




OBSERVED WIND 200MB FEB 1979

SCALE = 20




OBSERVED WIND 200MB FEB 1979


SCALE = 20




STREAM FUNCTION 200MB FEB 1979




STREAM FUNCTION 200MB FEB 1979



VELOCITY POTENTIAL 200MB FEB 1979




VELOCITY POTENTIAL 200MB FEB 1979




SCALE = 20

OBSERVED WIND 200MB NOV 1979




SCALE = 20


OBSERVED WIND 200MB NOV 1979




STREAM FUNCTION 200MB NOV 1979




STREAM FUNCTION 200MB NOV 1979





VELOCITY POTENTIAL 200MB NOV 1978




VELOCITY POTENTIAL 200MB NOV 1978







STREAM FUNCTION 200MB DEC 1978




STREAM FUNCTION 200MB DEC 1978



VELOCITY POTENTIAL 200MB DEC 1979




VELOCITY POTENTIAL 200MB DEC 1979




OBSERVED WIND 200MB JAN 1980

SCALE = 20




OBSERVED WIND 200MB JAN 1980


SCALE = 20




STREAM FUNCTION 200MB JAN 1980




STREAM FUNCTION 200MB JAN 1980



VELOCITY POTENTIAL 200MB JAN 1980




VELOCITY POTENTIAL 200MB JAN 1980




OBSERVED WIND 200MB FEB 1980

SCALE = 20




OBSERVED WIND 200MB FEB 1980


SCALE = 20




STREAM FUNCTION 200MB FEB 1980




STREAM FUNCTION 200MB FEB 1980



VELOCITY POTENTIAL 200MB FEB 1980



VELOCITY POTENTIAL 200MB FEB 1980




OBSERVED WIND 200MB NOV 1980

SCALE = 20




OBSERVED WIND 200MB NOV 1980


SCALE = 20




STREAM FUNCTION 200MB NOV 1980




STREAM FUNCTION 200MB NOV 1980



VELOCITY POTENTIAL 200MB NOV 1980




VELOCITY POTENTIAL 200MB NOV 1980




OBSERVED WIND 200MB DEC 1980

SCALE = 20




OBSERVED WIND 200MB DEC 1980


SCALE = 20




STREAM FUNCTION 200MB DEC 1980




STREAM FUNCTION 200MB DEC 1980



VELOCITY POTENTIAL 200MB DEC 1980




VELOCITY POTENTIAL 200MB DEC 1980




OBSERVED WIND 200MB JAN 1981

SCALE = 20




OBSERVED WIND 200MB JAN 1981


SCALE = 20




STREAM FUNCTION 200MB JAN 1981




STREAM FUNCTION 200MB JAN 1981



VELOCITY POTENTIAL 200MB JAN 1981



VELOCITY POTENTIAL 200MB JAN 1981




OBSERVED WIND 200MB FEB 1981

SCALE = 20




OBSERVED WIND 200MB FEB 1981


SCALE = 20




STREAM FUNCTION 200MB FEB 1981




STREAM FUNCTION 200MB FEB 1981



VELOCITY POTENTIAL 200MB FEB 1981




VELOCITY POTENTIAL 200MB FEB 1981




OBSERVED WIND 200MB NOV 1981

SCALE = 20




OBSERVED WIND 200MB NOV 1981


SCALE = 20




STREAM FUNCTION 200MB NOV 1981




STREAM FUNCTION 200MB NOV 1981



VELOCITY POTENTIAL 200MB NOV 1981




VELOCITY POTENTIAL 200MB NOV 1981




SCALE = 20

OBSERVED WIND 200MB DEC 1981




SCALE = 20


OBSERVED WIND 200MB DEC 1981



VELOCITY POTENTIAL 200MB DEC 1981



VELOCITY POTENTIAL 200MB DEC 1981




SCALE = 20

OBSERVED WIND 200MB JAN 1982




SCALE = 20


OBSERVED WIND 200MB JAN 1982




STREAM FUNCTION 200MB JAN 1982




STREAM FUNCTION 200MB JAN 1982



VELOCITY POTENTIAL 200MB JAN 1982




VELOCITY POTENTIAL 200MB JAN 1982




OBSERVED WIND 200MB FEB 1982

SCALE = 20




OBSERVED WIND 200MB FEB 1982


SCALE = 20




STREAM FUNCTION 200MB FEB 1982




STREAM FUNCTION 200MB FEB 1982



VELOCITY POTENTIAL 200MB FEB 1982



VELOCITY POTENTIAL 200MB FEB 1982




OBSERVED WIND 200MB NOV 1982

SCALE = 20




OBSERVED WIND 200MB NOV 1982


SCALE = 20




STREAM FUNCTION 200MB NOV 1982




STREAM FUNCTION 200MB NOV 1982



VELOCITY POTENTIAL 200MB NOV 1982



VELOCITY POTENTIAL 200MB NOV 1982




OBSERVED WIND 200MB DEC 1982

SCALE = 20




OBSERVED WIND 200MB DEC 1982


SCALE = 20




STREAM FUNCTION 200MB DEC 1982




STREAM FUNCTION 200MB DEC 1982



VELOCITY POTENTIAL 200MB DEC 1982



VELOCITY POTENTIAL 200MB DEC 1982




OBSERVED WIND 200MB JAN 1983

SCALE = 20




OBSERVED WIND 200MB JAN 1983


SCALE = 20




STREAM FUNCTION 200MB JAN 1983




STREAM FUNCTION 200MB JAN 1983



VELOCITY POTENTIAL 200MB JAN 1983




VELOCITY POTENTIAL 200MB JAN 1983




OBSERVED WIND 200MB FEB 1983

SCALE = 20




OBSERVED WIND 200MB FEB 1983


SCALE = 20




STREAM FUNCTION 200MB FEB 1983




STREAM FUNCTION 200MB FEB 1983



VELOCITY POTENTIAL 200MB FEB 1983



VELOCITY POTENTIAL 200MB FEB 1983



DISTRIBUTION LIST

No. Copies

1. Defense Technical Information Center
Cameron Station
Alexandria, Virginia 22314
2. Library, Code 0142
Naval Postgraduate School
Monterey, California 93943
3. Department of Meteorology Library
Code 63, Naval Postgraduate School
Monterey, California 93943
4. Climate Analysis Center
NMC/NOAA
Washington, D.C. 20233
5. Professor C.-P. Chang, Code 63Cp
Naval Postgraduate School
Monterey, California 93943
6. Professor J. Boyle, Code 63Xj
Naval Postgraduate School
Monterey, California 93943
7. Dr. K. M. Lau
Goddard Space Flight Center
Climate and Radiation Branch
NASA, Code 915
Greenbelt, Maryland 20771

2

2

1

2

40

2

2

END

FILMED

11-84

DTIC